

AD-A076 162 ARMED FORCES FOOD SCIENCE ESTABLISHMENT SCOTTSDALE (A--ETC F/G 6/8  
HUMAN DIETARY FIBRE: A REVIEW, (U)  
MAY 79 C H FORBES-EWAN

UNCLASSIFIED AFFSE-4/79

NL

| OF |  
AD  
A076 162



END  
DATE  
FILMED

11-79

DDC

UNCLASSIFIED

B. 5. 12

AFFSE REPORT 4/79

AR No. 001-711

AD A 076162



LEVEL #



Department of Defence  
Defence Science and Technology Organisation  
Armed Forces Food Science Establishment  
Scottsdale, Tasmania

AFFSE REPORT 4/79

© COMMONWEALTH OF AUSTRALIA, 1979

DDC FILE COPY

## Human Dietary Fibre : A Review



C. H. FORBES-EWAN

79 11 05 053 May 1979

THE UNITED STATES NATIONAL  
TECHNICAL INFORMATION SERVICE  
IS AUTHORIZED TO  
REPRODUCE AND SELL THIS REPORT

APPROVED  
FOR PUBLIC RELEASE

UNCLASSIFIED

DEPARTMENT OF DEFENCE

*Armed Forces Food Science Establishment*

14 AFFSE REPORT - 4/79

6 HUMAN DIETARY FIBRE: A REVIEW (U)

10 C. H. FORBES-EWAN

11 May 79 12 21

© Commonwealth of Australia, 1979

SUMMARY

A review is made of the composition, analysis and relationship to disease of fibre in human nutrition. (U)

---

POSTAL ADDRESS: The Director,  
Armed Forces Food Science Establishment,  
P.O. Box 147,  
Scottsdale, Tasmania, 7254.

---

410 231 int



# DOCUMENT CONTROL DATA SHEET

UNCLAS

- |                                |                                    |
|--------------------------------|------------------------------------|
| 1. DOCUMENT NUMBERS            | 2. SECURITY CLASSIFICATION         |
| a. AR Number: 001-711          | a. Complete document:<br>Unclas    |
| b. Document Series and Number: | b. Title in isolation:<br>Unclas   |
| c. Report Number: 4/79         | c. Summary in isolation:<br>Unclas |

3. TITLE: Human Dietary Fibre: A Review

- |  |                                |
|--|--------------------------------|
| 4. PERSONAL AUTHORS:<br>Forbes-Ewan, C. H. | 5. DOCUMENT DATE:<br>May, 1979 |
|--|--------------------------------|

6. TYPE OF REPORT AND  
PERIOD COVERED:  
Technical Report

- |  |   |
|--|---|
| 7. CORPORATE AUTHOR:<br>Armed Forces Food Science<br>Establishment,<br>Scottsdale, Tasmania, Aust. | 8. REFERENCE NUMBERS:<br>a. Task: FSE 76/226<br>b. Sponsoring Agency:<br>DOD (Army) |
|--|---|

9. COST CODE: 241

- |                                   |                       |
|-----------------------------------|-----------------------|
| 10. IMPRINT:<br>AFFSE — May, 1979 | 11. COMPUTER PROGRAM: |
|-----------------------------------|-----------------------|

12. RELEASE LIMITATIONS:  
Approved for public release

12-0 OVERSEAS: N.O. ☐ P.R. ☐ 1 ☐ A ☐ B ☐ C ☐ D ☐ E ☐

13. ANNOUNCEMENT LIMITATIONS: —

- |   |                                |
|---|--------------------------------|
| 14. DESCRIPTORS:<br>Nutrition, chemical analysis, fibre | 15. COSATI CODES:<br>0608-0616 |
|---|--------------------------------|

16. SUMMARY:  
A review is made of the composition, analysis, and relationship to  
disease of fibre in human nutrition. (U)

UNCLAS

## CONTENTS

	Page No.
Definition of Fibre .....	1
Components of Fibre and their possible Physiological Significance .....	1
Arrangement and Distribution of the Components of Dietary Fibre in Foods	4
The Analysis of Dietary Fibre .....	5
Possible Modes of Action .....	6
Theories on the Health-Benefiting Role of Fibre in Diets .....	6
Dietary Fibre and Lipid Metabolism .....	7
Dietary Fibre and Colon Function .....	8
Epidemiology of Bowel Disease .....	8
Dietary Fibre and Diverticular Disease .....	9
Dietary Fibre and Colonic Cancer .....	10
Fibre and Appendicitis .....	11
Fibre and Dental Caries .....	11
References	
Distribution List	

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Availand/or special
A	

# HUMAN DIETARY FIBRE

## A REVIEW OF THE COMPOSITION, ANALYSIS AND RELATIONSHIP TO DISEASE OF FIBRE IN HUMAN NUTRITION

### Definition of Fibre

The term "fibre" is at present an imprecise term. The concept arose initially to denote that component of a forage crop which was unavailable to animals. This was referred to as "crude fibre".

Crude fibre analysis consists of extraction with petroleum ether, followed by acid, then alkali digestion (i.e. simulating animal digestion). The undigested residue is weighed, ashed and the crude fibre value is the final residue weight minus the ash weight (Horwitz, 1970).

The concept of "fibre", particularly in relation to human nutrition has altered over the years, until now "fibre" is considered to be the whole of the plant cell wall plus some substances such as mucilages and gums which are not found in the cell wall but which are indigestible in the human gut. Not all plant cell wall constituents are indigestible though, so fibre is not completely indigestible, and not all the constituents of "dietary fibre" are, in fact, fibrous. (Cummings, 1973).

The crude fibre analysis described above is extremely method-dependent and insensitive at the low fibre levels found in many foods (van Soest, 1963). It also measures a very variable proportion of the total plant cell wall constituents, which are invariably underestimated. Crude fibre analysis recovers 50-80% of the cellulose, 10-50% of the lignin and only about 20% of the hemicelluloses (van Soest, 1973). The proportion of the total cell wall constituents recovered depends to a large extent on the percentage of hemicellulose present. Hemicelluloses are largely lost in the crude fibre analysis.

The crude fibre method thus has many deficiencies and does not have a place in the analysis of dietary fibre in human foods.

Alternative methods of analysis will be discussed later.

### Components of Fibre and their Possible Physiological Significance

- (1) **Cellulose.** This is the only truly "fibrous" component of the cell wall. This is an unbranched 1-4B-D-glucose polymer containing about 3000 glucose units. The whole molecule folds into a flat, ribbon-like structure which packs closely with identical molecules to form micro-fibrils. These form the fibres of cellulose which are woven into the plant cell wall (Ranby, 1969).

The fibres are very strong due, it is believed, to the length of the chains and hydrogen bonding between hydroxyl groups of adjoining polymers (Jeffrey

and Rosenstein, 1964). For the most part, cellulose is crystalline, but there are regions along the micro-fibrils which are non-crystalline (amorphous).

In terms of significance to human nutrition, the most important properties of cellulose are the susceptibility of the molecule to enzymatic and acid hydrolysis, and its ability to absorb water. Both these characteristics appear to involve the amorphous areas of the microfibrils. Cellulose is susceptible to acid hydrolysis and is 30-40% digested (Mangold, 1934).

Most plant cell walls contain 15-40% pure cellulose on a dry-weight basis (Kent-Jones and Amos, 1967).

Inclusion of cellulose in the diet of both animals and man is associated with changes in cholesterol and bile acid metabolism, however cellulose appears to be less potent than other dietary fibre constituents in lowering blood cholesterol level (Keys et al, 1961).

- (2) **Hemicelluloses.** Mostly 5-carbon sugars, they are soluble in cold, dilute alkali. Hemicelluloses are either acidic or neutral depending on whether they have small or large numbers of uronic acid residues. Together with pectin, the hemicelluloses form the matrix of the plant cell wall in which are enmeshed the cellulose fibres.

Of possible physiological significance are: hemicelluloses have a water-holding capacity, are 55-90% digestible (Williams and Olmsted, 1936, Southgate and Durnin, 1970) and bind ions.

- (3) **Pectins.** Pectins constitute between 1 and 4% of total cell wall polysaccharides (Whistler and Richards, 1970). The parent molecule is a polymer of 1-4-B-D-galacturonic acid. The parent polymer is usually modified to include 10-25% of neutral sugar residues. Pectins have the ability to form gels and can bind ions. In man, pectins may be almost completely digested (Wersh and Ivy, 1941). Pectins can alter cholesterol metabolism (lowering blood cholesterol level). In man, pectin supplements of 15-35 g per day led to significant lowering of blood cholesterol (Fisher et al, 1965).

In another experiment, pectin supplements were found to be second only to a reduction in dietary intake of cholesterol in reducing blood cholesterol levels (Anderson, 1973).

The degree of methylation of the molecules affects both the degree of gel-formation and the cholesterol lowering ability (Ershoff and Wells, 1962).

- (4) **Plant gums, mucilages, storage polysaccharides.** These are not cell-wall components, but are related to them biochemically and in their physiological effects. Mucilages are found in seeds where they protect against desiccation by holding water. Mostly they are neutral polysaccharides.

Structurally, they resemble hemicelluloses. Guar is an example; in small amounts it is used as a food thickener.

Gums form at the sites of injury in plants. In the food industry they are used as stabilizers, emulsifiers, thickeners and in bulk-forming laxatives.



Gum arabic is also used as an adhesive.

Gums are a complex group of highly branched uronic-acid-containing polymers, with neutral sugars such as xylose and arabinose. A high proportion of calcium and magnesium salts is formed and a significant number of residues are acetylated (Aspinall, 1970).

In terms of physiological effects on man, the most important characteristics of gums and mucilages are their effects on cholesterol and bile salt metabolism. They have similar effects to pectin. Guar is especially potent (Jenkins et al, 1975).

- (5) **Lignin.** This is the atypical constituent of fibre in that it is not a carbohydrate. It is a small, insoluble polymer with molecular weight 1000-4000. It is made up of three different subunits, each of which consists of a substituted phenylpropane (Kent-Jones and Amos, 1967).

Lignin's role in the cell wall is to strengthen it by permeating and supporting the other components (Pearl, 1967). Lignin impairs the digestibility of other cell wall constituents (Raymond, 1969). It is also thought to have a bile-salt-binding capacity, with adsorption perhaps occurring through hydrophobic interactions (Eastwood and Hamilton, 1968).

- (6) **Algal Polysaccharides.** Related to the above components of fibre is alginic acid, derived from algae. This is a nonsulphated polysaccharide. In its native form it occurs as sodium, calcium, potassium or magnesium salts (alginates). It is a linear, acid, polysaccharide polymer, largely unbranched so it, like cellulose, can form microfibrils.

Alginic acid is widely used in the food, pharmaceutical and paper industries. In food, it impairs the absorption of strontium in the gut (Patrick, 1967).

- (7) **Fibre-associated substances.** Methods of fibre analysis invariably leave some non-fibre substances in the residue. The extent to which some of the characteristics of fibre are in fact due to these contaminants is not known.

Phytic acid is found in seeds in association with protein. Usually it is complexed with Ca, Mg and K and exists as salts (phytin). Phytin is milled off with the bran during the refining of wheaten flour; it is therefore present in wholemeal bread and not, significantly, in white bread (Kent-Jones, 1967). Because of the ion-binding capacity of phytin, the consumption of wholemeal bread can deplete ions such as calcium from the gut. In areas where calcium intake is borderline, this could lead to osteomalacia (Berlyne et al, 1973).

Phytin is also thought to impair iron and zinc absorption (Jenkins et al, 1976). The principal constituent of the ash in the fibre fraction is silicon. Its significance to human nutrition relates to its capacity to impair digestibility of cell-wall materials. In this respect it bears some similarity to lignin (Raymond, 1969).

Cuticular substances (waxes and cutins) also apparently impair cell-wall digestion (van Soest and McQueen, 1973).

The plant cell wall contains a multitude of other substances whose role in human nutrition is not clear, e.g. tannins, enzymes, vitamins, glycosides, etc.

**Conclusion.** Fibre is a complex matrix of polymers from the plant cell-wall and similar substances from elsewhere in the plant cell, e.g. gums and mucilages.

The main physiological effects of these individual components are on: cholesterol and bile acid metabolism, water-holding capacity, gel-formation, and ion-binding.

The physical and chemical interactions of the constituents of fibre are important in the context of nutrition, as the components do not exist separately. How the physiological effects of individual components are altered by the presence of other fibre constituents is unknown.

### **Arrangement and Distribution of the Components of Dietary Fibre in Foods**

- (1) **Development of the cell wall.** (Albersheim, 1965). After nuclear division of the parent cell, a cell plate forms between the daughter nuclei. This is rich in pectic substances and becomes the middle lamella. The primary wall, which is deposited on the middle lamella, consists of cellulose fibrils that appear to be deposited as a random network. The matrix that forms around these fibrils contains non-cellulosic polysaccharides with both pectic and hemicellulosic characteristics.

The secondary wall usually consists of three distinct layers. Cellulose fibrils are laid down in a regular parallel arrangement. The matrix surrounding these is primarily hemicellulosic.

Lignification usually starts in the middle lamella and then proceeds outwards, swelling the cell wall. The actual composition of the cell wall varies from tissue to tissue and depends on the age of the plant (particularly with respect to lignin). Hirst, et al, 1959):

- (i) **Stem structures:** Most "stemfoods" are consumed in immature condition, are lightly lignified with a cuticular layer on the outer surface. Potato is an example.
- (ii) **Leaf Structure:** These are lightly lignified, have a cuticular layer and a waxy coating. In monocotyledons, heavy lignification occurs.
- (iii) **Root Structure:** These are usually eaten before much secondary thickening occurs. They are therefore lightly lignified, e.g. carrots, turnips.
- (iv) **Fruits:** These vary widely. There are two distinct classes, fleshy (e.g. tomato, citrus) and hard (e.g. nuts). Fleshy fruits are typically thin-walled, with only light lignification.
- (v) **Seeds.** The walls of the cells in the seed coat are often thickened and frequently lignified. In some, mucilages are present which absorb water and swell, thus rupturing the seed coat and cuticle (i.e. germinating).



## The Analysis of Dietary Fibre

As mentioned in Section (a), the "crude fibre" analytical method is not applicable to human dietary fibre with any degree of precision.

As yet, no one method is available which can be used on all foods without modification.

The "acid detergent fibre" (ADF) method of van Soest (1963) provides an estimate of fibre that correlates closely with the nutritive value of ruminant feedstuffs. The fraction measured by this procedure has been shown to consist mainly of cellulose and lignin. It also contains some silicon.

The method has a number of limitations when applied to human foods (Southgate, 1976). Human foods are usually much higher in lipids and starch than animal forage crops; this leads to problems of foaming and filtration. A more fundamental problem is that the ADF method measures only a fraction, albeit a significant one, of the components of human dietary fibre.

The "neutral detergent fibre" (NDF) method also of van Soest (1963) is thought to give an accurate measure of the cell wall constituents of vegetables

According to Southgate (1976) it appears to divide the dry matter of feeds very nearly into those constituents which are nutritionally available to man and those which depend on microbial fermentation.

The method involves extraction of non-fibre components of food in a neutral detergent followed by filtration and ashing of the residue to correct for inorganic contamination of the fibre.

The main problems (Southgate, 1976) are:

- (1) Imprecision due to small sample size. If a larger sample is used, problems of foaming occur. This can be largely overcome by initial lipid extraction.
- (2) Starch can remain insoluble in the hot detergent and be measured with residual NDF. Enzymatic hydrolysis can correct for this.
- (3) A fundamental problem is that the method will underestimate total dietary fibre because the water soluble polysaccharides are lost during the extraction. Angus et al (1977) on the other hand, claim that the NDF method leaves as fibre appreciable quantities of material susceptible to digestion by mammalian digestive enzymes.

They propose a method based on the NDF technique which includes water, ethanol and ether extraction and pancreatin digestion, as well as the neutral detergent extraction.

Their results were consistently lower (i.e. they obtained less fibre) than those obtained using the NDF method.

McCance et al (1936) measured "unavailable carbohydrate", a fraction which corresponds closely to dietary fibre, in a range of fruits, nuts and vegetables. They determined the residue insoluble in 80% v/v ethanol and then measured the starch in the residue after enzymatic hydrolysis with a takadiastase preparation; they then corrected for protein in the residue.

Southgate and Durnin (1970) developed methods to measure the various components of the unavailable carbohydrate fraction and thus form a basis for the measurement of dietary fibre. The method is, of necessity, very time-consuming and could not be used for routine analytical work. The first part of the method, however, gives a similar estimate of total dietary fibre to the method of McCance et al (1936). Southgate's method is based on an alcohol-insoluble residue, with starch removed by enzymatic hydrolysis and fat removed by diethyl ether extraction.

**Conclusions:** No method of analysis for dietary fibre has yet reached the stage of an accepted, standard method, applicable to all, or a majority of human foods. Modifications of the NDF method and the initial stages of the Southgate method seem to offer the best promise of a routine method.

### Possible Modes of Action

Vegetable fibre is a physical complex that acts principally in the colon. Fibre has water-holding, cation exchange and adsorptive properties, which vary with the variety of fruits and vegetables studied. It is possible that vegetable dietary fibre acts in the colon by forming a supporting matrix which provides surfaces or a locating micro-environment in which bacteria and intestinal contents interact (Eastwood and Mitchell, 1976). Vegetable dietary fibre may also act as a gel with partitioning characteristics for both bacteria and other substances in the intestinal contents. The metabolism of substances in the colon in its turn is influenced by the physical characteristics of the matrix, and this supporting medium may be metabolized and its character changed by the activity of the bacteria. There are important effects of fibre on solute and bacterial activity, which influence the fate of bacterial metabolites of solutes of endogenous and exogenous origins. Such effects may decide whether the metabolites are excreted in the faeces or are conserved by being returned in the portal vein. Fibre itself may be metabolized by bacteria (Wersh and Ivy, 1941).

It is possible to protect against the carcinogenic effect of chemical carcinogens by inducing increased mixed-function oxidase activity. The level of mixed function oxidase activity in the small intestine is related to the amount of fibre in the diet. The factor(s) is not known, but it could be lignin which does act as an antioxidant in the digestive tract (Wattenberg, 1974).

The significance to the human host of the colonic microbial activity is not clear. The acids formed in the posterior gut by fermentation of protein may have value, and vitamin synthesis may be important, but many nitrogenous products of dubious or negative value are also formed, in contrast to the fermentation products from carbohydrates (i.e. in herbivores such as ruminants) which give rise to volatile fatty acids which are absorbed by the host and oxidized to provide energy (Hungate, 1976).

### Theories on the Health-Benefiting Role of Fibre in Diets

- (1) Fibre functions as a Bulking Agent: Undigested, high molecular weight compounds form faecal bulk. Bulk in the intestinal tract induces a faster transit time and reduces the opportunity for microbial production of toxins or carcinogens. It has been suggested that cancer of the bowel can be due to the production of carcinogens by the action of colonic bacteria on compounds present in the colon, e.g. bile acids and steroids (Burkitt, 1974).

- (2) Fibre functions as a toxin antagonist: Large, insoluble compounds are ideal absorbants. Dietary fibre could bind potential toxins, prevent their digestion or absorption and facilitate their excretion.
- (3) Fibre Functions as a Nutrient: Fibre may be a source of unknown nutrients or compounds that have medicinal properties. Natural plant fibres are a heterogeneous mixture of complex compounds. The chemical structure and biological activity of the individual molecule is often unknown. Even undigested molecules can cause an indirect health benefit by stimulating the growth of beneficial microflora or by retarding the growth of pathogenic organisms (Lang and Briggs, 1974).

**Animal Studies.** In monogastric animals, e.g. rat, mouse, rabbit, chick and guinea pig, fibre seems to be non-essential, except perhaps in the guinea pig, although it confers benefits to the rat and rabbit and to many other species (Lang and Briggs, 1976).

**Conclusion.** The debate over the need for dietary fibre has a long history and will undoubtedly continue for many decades. Currently, we must conclude that we have not seen convincing evidence that dietary fibre is an essential dietary component. The development of chemically defined liquid diets for rats, chickens and man suggests that there is no unique, essential function for dietary fibre.

Although dietary fibre may not be an essential component, many studies indicate that it may have health-benefiting properties, such as reducing the effects of toxins, reducing cholesterol levels and preventing diverticular disease. Unfortunately, many of these theories involve slow processes and thus require long-term tests which must be carefully controlled.

Excessive fibre may be harmful, e.g. phytic acid, oxalic acid and phenolic compounds are associated with plant fibres and have been shown to restrict the availability of essential elements such as zinc (National Research Council, 1973).

In the context of possible benefits conferred by dietary fibre, it is interesting to note that placebos frequently consist of fibrous material, such as carboxymethylcellulose. A consistent proportion of patients has been cured by placebo treatment, a result which is usually attributed to psychological factors. In view of the antitoxic effects of cellulose, perhaps placebos should not contain fibre (Lang and Briggs, 1976).

#### **Dietary Fibre and Lipid Metabolism (Danielsson, 1973)**

Cholesterol is a monohydric sterol in every cell in the body. It is synthesized mainly in the liver and intestine. The major catabolic products are the bile acids. The bile contains cholesterol, bile salts and phospholipids (primarily lecithin) in a delicate balance. Any substantial change in any one of these can lead to gallstones.

In general, pectin and other polysaccharides are hypo-cholesterolaemic (Kirayama et al, 1969). The suggestion has been made that the mechanism of action of pectin is inhibition of cholesterol absorption and increase of faecal bile acid excretion (Leveille and Sauberlich, 1966). Experiments involving man have not shown a great effect of bran on serum cholesterol level (Connel et



al, 1975). Pectin and guar gum, on the other hand seem to be quite effective in lowering serum cholesterol (Keys et al, 1961). Evidence suggests that fibre may inhibit cholesterol absorption by binding bile salts. Such binding could result in a failure in micellar formation, essential for cholesterol absorption, and this, in turn, would increase bile acid excretion, effecting an increase in bile acid synthesis to replace the lost bile salts. Both events would drain cholesterol pools (Kritchevsky and Story, 1974).

Water-holding and cation-exchange properties of fibre may also influence absorption of cholesterol and bile acids (McConnell et al, 1974).

The accumulated data indicate a general hypolipidaemic and anti-atherogenic effect. However, the type of fibre is important, for example lignin actively binds bile salts in vitro, cellulose does not (Kritchevsky and Story, 1974).

### **Dietary Fibre and Colon Function**

- (1) Effect on Stool Weight: Fibre, especially from bran, leads to softer, heavier stools. The increase in weight appears to be due to an increase in water content of the stool (Eastwood et al, 1973). Fruits and vegetables as absorbers are: turnip < potato < banana < cauliflower < pea < winter cabbage < lettuce < apple < carrot < mango < bran (McConnell et al, 1974). Particle size is also important - coarse bran holds more water than fine bran (Kirwan et al, 1974).

The constituents of fibre responsible for the absorption of water have not been conclusively identified. Probably the polysaccharide content is involved.

The bulking action of dietary fibre will have the added effect of diluting colonic contents, e.g. bile acids, which according to Burkitt (1974) should lower the risk of colonic cancer.

- (2) Effect on transit time: Rural Africans who have a high fibre diet not only produce large, bulky stools, but are reported to have fast transit times, of the order of 30 hr, in contrast to Europeans and North Americans (< 48 hours) (Burkitt et al, 1972). Cereal bran, when given with the normal diet, not only causes an increase in stool weight, but also lowers transit times.

However, people with short transit times (<24 hours) react to bran by increasing their transit time (Harvey et al, 1973), perhaps suggesting that bran gives rise to an ideal "bran transit time", that will vary depending on the source of the vegetable fibre in the diet. The mechanism whereby fibre leads to a decreased transit time is not adequately explained. Vegetable fibre may alter the specific gravity of colonic contents; substances having a specific gravity of above or below 1.1 being more rapidly excreted than substances with specific gravity of 1.1 (Kirwan and Smith, 1974).

### **Epidemiology of Bowel Disease**

Non infective bowel diseases include haemorrhoids, appendicitis, diverticular disease, cancer of the colon, colitis and Crohn's disease. It is thought that a deficiency of fibre coincides with their appearance.

With some bowel diseases, an excess of dietary fibre appears to coincide: volvulus of the colon and inflammatory bowel diseases (Cummings, 1973).

Many countries, e.g. some African, parts of India and Japan (Burkitt et al, 1974) appear to be free of non-infective bowel disease. There is a temptation to ascribe these characteristics to race. However, when people from these areas migrate to a country with high incidence, they suffer the same rate of bowel disease as their adopted countrymen. This suggests an environmental cause (Burkitt, 1974a).

The reverse migration, people from high-incidence countries going to low-incidence areas, does not usually result in a decreased incidence of colonic disease (Burkitt, 1974a).

It is unlikely that people from affluent countries would become less prosperous on moving to poorer countries. Thus non-infective bowel disease appears to be an environmental disease, and the component of the environment concerned appears to be connected to affluence.

Strong arguments exist that food is the environmental factor involved (Burkitt, 1974). Africans in low incidence areas derive 80% of their food energy from high fibre carbohydrates, mainly yams, millet and sorghum (Latham, 1965). In the West, the tendency over the past century or so has been towards an increased consumption of white flour (low in fibre), coupled with a dramatic rise in white sugar consumption. Fat consumption has also increased in the West. There is a correlation between these changes in diet and the sharp increase in incidence of non-infective bowel diseases but the correlation is tenuous, due to lack of data on both changes in diet and changes in frequency of bowel diseases.

Cleave (1969) believes that increased consumption of sugar is the main factor. The evidence for this is not strong, consisting mainly of a correlation between increased sugar consumption and increased incidence of colonic disorders.

### **Dietary Fibre and Diverticular Disease**

This condition is common in Western countries but uncommon among Africans eating a high fibre diet (Painter and Burkitt, 1975).

It has been postulated that low fibre diets lead to the formation of small, hard stools which exert excessive pressure on the colon, causing small pockets or diverticula to be formed (Burkitt, 1974a).

It is estimated that a majority of people over the age of 60 years in Western countries have diverticula in the colon. However, only about 10% ever suffer symptoms (Painter and Burkitt, 1975).

Burkitt (1974a) suggests that bran should be added to the diet to protect against diverticular disease. Bran is the most concentrated form of fibre; it consists of about 21% cellulose, 20-26% pentosan (hemicellulose), 7.5-9.0% starch, 5% sugar, 11-15% protein, 5-10% fat, 5-9% ash and about 14% moisture (Kent-Jones and Amos, 1967).

That bran is beneficial in the treatment of diverticular disease is unquestionable. Many studies have shown improvement of patients after consuming

bran supplements (Burkitt, 1974a). It is highly likely that vegetable fibre will have a similar effect to that of bran; studies have not yet been carried out.

The problem of aetiology of diverticular disease is not solved by the finding that fibre will alleviate diverticular disease. It is possible that the cause of diverticulosis is something other than lack of fibre; there is no doubt, however, that fibre is of therapeutic value.

### **Dietary Fibre and Colonic Cancer**

There is widespread belief that diet and colonic cancer are related. The factor(s) that may be involved have not been established.

The low incidence of this cancer in Africa (e.g. in Nigeria, 6/100,000) compared with North American Negroes (70/100,000) suggests an environmental cause (Burkitt et al, 1972).

The rate of stomach cancer, in contrast, remains the same as in the country of origin (Drasar and Irving, 1973).

Burkitt (1974a) suggests that low levels of dietary fibre can lead to longer transit times and that bacteria in the colon can produce carcinogens from metabolites in the colon. The longer transit time would allow more of the carcinogens to be produced and would lead to a longer time of exposure to the carcinogens. Bile acids have been proposed as one substrate which bacteria may metabolise to produce carcinogenic by-products.

There are data to support the theory that carcinoma of the bowel is related to an enhanced faecal bile acid excretion (Hill et al, 1975). Cholesterol metabolism and colonic carcinoma may be inter-related. When there is an excess of serum cholesterol, it is converted to bile acids, which pass to the colon (Danielsson, 1973). Patients with large bowel cancer had higher faecal bile acid concentrations than the controls. More cancer patients also had nuclear dehydrogenating bacteria (Hill et al, 1975).

However, the correlation between low dietary fibre and cancer of the colon is weak.

Hill (1971) showed that the amount of bile acid in the faeces is dependent on the fat content of the diet. This may provoke a dilemma to heart patients who are urged to take polyunsaturated fats which lower the serum cholesterol level, but lead to increased faecal bile acid excretion.

Thwe et al (1976) found no significant difference in bacterial flora and daily excretion of bile acids in fast transit (<24 hrs) and slow transit (<48 hrs) people.

Because of the water-holding capacity of fibre, stools formed by people on high fibre diets tend to be much heavier (about 500 g) than those formed as a result of low-fibre diets. Thus Findlay et al (1974) showed that when the diet was supplemented with 20g bran per day, stool weight increased, but the concentration of bile acids decreased. This dilution of potential carcinogens may be of significance in reducing the risk of bowel cancer.

There is, however, a closer correlation between high fat and protein diets and cancer of the colon than between low fibre and bowel cancer (Wynder and



Shigematsu, 1967). Among potential carcinogens are several amino acids and lecithin, which may be converted to nitrosamine. N-nitroso compounds present in bacon, smoked fish and mushrooms may be converted to carcinogens (Lancet, 1973). Reducing agents, such as ascorbic acid, may have an unremarked protective effect by preventing the conversion of potential carcinogens to actual carcinogens.

The problem of carcinogenesis probably devolves on bacterial activity, to some extent, determining whether potential carcinogens are converted to actual carcinogens. Fibre will probably be important because of its dilution effects.

### **Fibre and Appendicitis**

Studies in South Africa have shown a strong correlation between a diet high in fibre and low rates of appendectomy (Walker, A.R.P. et al, 1973). Among students in various parts of South Africa, rural Negroes had a rate of 0.5% appendectomy, Whites had a rate of 16.5%. In the cities, Negroes and Whites had similar appendectomy rates. Rural Negroes consume far greater amounts of vegetable fibre than either city Negroes or Whites.

Burkitt (1974) believes that a fibre-depleted diet can result in the formation of fecaliths and also excessive segmentation of the appendix, either of which may obstruct the appendix lumen. Obstruction might raise the intraluminal pressure sufficiently to devitalize the appendicular mucosa and allow bacterial invasion.

The evidence indicates that a change from a refined diet to one less refined is accompanied by a reduced incidence of the disease.

### **Fibre and Dental Caries**

The common acceptance of the aetiology of dental caries as high sugar consumption is not supported by all the available evidence.

On the Island of Lewis (Scotland), white children consuming 100 g sugar per day had an average of about 2 decayed, missing or filled teeth (DMF) (King, 1940). White children in Johannesburg eating similar amounts of sugar had DMF of 10 (Retief et al, 1975).

Other evidence points to an increase in fibre consumption, as perhaps being of importance in the alleviation of dental caries. During World War II, the caries situation in children improved in all European countries (Bowen, 1972). The change was associated with a fall in sugar intake and a rise in consumption of fibre contained in cereals, vegetables and fruits. However, in Germany, dental caries also decreased, but this was in spite of a rise in sugar consumption (Wandett, 1969). This suggests that alterations other than a decrease in sugar intake were instrumental in effecting a reduction in dental caries. As with Western Europe, the diet increased in fibre in Germany during the war.

### **Conclusion**

The "Fibre hypothesis" at present is more of an association than a relationship; that is, correlations have been shown to exist between various diseases and a decrease in dietary fibre.

Fibre undoubtedly has a therapeutic value in diverticulosis and at least eases the discomfort of haemorrhoids. Whether a fibre deficiency is involved in the aetiology of all or some of the diseases discussed above is, as yet, unproved.

## REFERENCES

- Albersheim, P. (1965). Biogenesis of the cell wall. In: Plant Biochemistry. J. Bonner and J. E. Varner (eds.) Academic Press.
- Anderson, J. T. F. et al (1973). *J. Am. Diet. Associ* **62**: 133. Cholesterol-lowering diets.
- Angus, R. et al (1977). *Proc. Nutr. Soc. Aust.* **2**: 90. A simplified method for determining fibre in food.
- Aspinall, G. O. (1970). Pectins, plant gums and other plant polysaccharides. In: The carbohydrates, W. Pigman and D. Horton (eds.) New York, Academic Press, Vol IIB, Chap 39, p. 515.
- Bowen, W. H. (1972). *Arch. Dis. Child.* **47**: 849. Dental Caries.
- Burkitt, D. P. (1971). *Br. J. Surg.* **58**: 695. The aetiology of appendicitis.
- Burkitt, D. P. et al (1972). *Lancet* **2**: 1408. Effect of dietary fibre on stools and transit times and its roles in the causation of disease.
- Burkitt, D. P. et al (1972). *Dis. Colon. Rectum* **17**: 456. An epidemiologist approach to cancer of the large intestine.
- Burkitt, D. P. et al (1974a). *J.A.M.A.* **229**: 1068. Dietary fibre and disease.
- Burlyne, G. M. et al (1973). *Am. J. Clin. Nutr.* **26**: 910.  
Bedouin osteomalacia due to calcium deprivation caused by high phytic acid content of unleavened bread.
- Cleave, J. L. et al (1969). Diabetes, Coronary Thrombosis and the Saccharine Disease. 2nd ed., Bristol, J. Wright and Sons.
- Connell, A. M. et al (1975). *Lancet* **1**: 496. Absence of effect of bran on blood lipids.
- Cummings, J. H. (1973). *Gut* **14**: 69. Dietary Fibre.
- Danielsson, H. (1973). Mechanism of bile-acid biosynthesis. In: The Bile Acids, Vol 2, P. Nair and D. Kritchevsky (eds). Plenum, p. 1.
- Drasar, B. S. and D. Irving (1973). *Br. J. Cancer* **27**: 167. Environmental factors and cancer of the colon and breast.
- Eastwood, M. A. and D. Hamilton (1968). *Biochem. Biophys. Acta* **152**: 165. Studies on the adsorption of bile salts to non-absorbed components of the diet.
- Eastwood, M. A. et al (1973). *Br. Med. J.* **4**: 392. Effects of dietary supplements of wheat bran and cellulose on faeces and bowel function.
- Eastwood, M. A. and Mitchell, W. D. (1976). In: Fibre in Human Nutrition. G. Spiller and R. Amen (eds.) Plenum p. 123.
- Findlay, J. M. et al (1974). *Lancet* **1**: 146. Effects of unprocessed bran on colon function in normal subjects and in diverticular disease.
- Fisher, H. et al (1965). *J. Nutr.* **86**: 113. Dietary pectin and blood cholesterol.
- Harvey, R. F. et al (1973). *Lancet* **1**: 1278. Effects of increased dietary fibre on intestinal transit.
- Hill, M. (1971). *J. Pathol.* **104**: 239. The effect of some factors on the faecal concentration of acid steroids, neutral steroids and urobilins.
- Hill, M. et al (1975). *Lancet* **11**: 535. Faecal bile acids and clostridia in patients with cancer of the large bowel.

- Hirst, E. L. et al (1959). *J. Sci. Food Agric.* **10**: 19. Analytical studies on the carbohydrates of grasses and clovers. IX. Changes in carbohydrate composition during the growth of lucerne.
- Horwitz, W. (1970). Crude Fibre. In: *Official Methods of Analysis of Official Agricultural Chemists, U.S.A.*, 11th ed p. 129.
- Hungate, R. E. (1976). In: *Fibre in Human Nutrition*. G. Spiller and R. Amen (eds). Plenum. p. 148.
- Jeffrey, G. A. and R. D. Rosenstein (1964). *Adv. Carbohydr. Chem.* **19**: 7. Crystal structure analysis in carbohydrate chemistry.
- Jenkins, D. J. A. et al (1975). *Lancet* **1**: 1116. Effect of pectin, guar gum and wheat fibre on serum cholesterol.
- Jenkins, D. J. A. et al (1976). *Am. J. Clin. Nutr.* **28**: 1408. Effect of wheat fibre on blood lipids, faecal steroid excretion and serum iron.
- Kent-Jones, D. W. and A. J. Amos (1967). *Modern Cereal Chemistry*, 6th Ed. London, Food Trade Press.
- Keys, A. et al (1961). *Proc. Soc. Exp. Biol. Med.* **106**: 555. Fibre and pectin in the diet and serum cholesterol concentration in man.
- King, J. D. (1940). *Med. Res. Counc. Spec. Rep. Ser. No. 241*. London H.M. Stat. Office. Dental disease in the Island of Lewis.
- Kirwan, W. O. et al (1974). *Br. Med. J.* **2**: 187. Action of different bran preparations on colonic function.
- Kirwan, W. O. and A. N. Smith (1974). *Sc. J. Gastroenterol* **9**: 763. Gastrointestinal transit time estimated by an isotope capsule.
- Kiriyama, S. et al (1969). *J. Nutr.* **97**: 382. Hypocholesterolaemic effect of polysaccharides and polysaccharide - rich foodstuffs in cholesterol-fed rats.
- Kritchevsky, D. and J. A. Story (1974). *J. Nutr.* **104**: 458. Binding of bile salts in vitro by non nutritive fibre.
- Lancet* (1973). *Leader* **2**: 1243. Environmental nitrosamines.
- Lang, J. A. and G. M. Briggs (1976). In *Fibre in Human Nutrition*. G. Spiller and R. Amen (eds). Plenum pp 162-166.
- Latham, M. C. (1965). *Human Nutrition in Tropical Africa*. Rome, Italy, F.A.O.
- Leveille, G. A. and Sauberlich, H. E. (1966). *J. Nutr.* **88**: 209. Mechanism of the cholesterol-depressing effect of pectin in the cholesterol fed rat.
- Mangold, D. E. (1934). *Nutr. Abstr. Rev.* **3**: 647. The digestion and utilization of crude fibre.
- McCance, R. A. et al (1936). *Med. Res. Counc. Spec. Rep. Ser.* 213 (HMSO): 32. The nutritive value of fruits, vegetables and nuts.
- McConnell, A. A. et al (1974). *J. Sci. Food Agric.* **25**: 1457. Physical characteristics of vegetable foodstuffs that could influence bowel function.
- National Research Council (1973). *Toxicants occurring naturally in foods*. Washington, D.C. National Academy of Sciences.
- Painter, N. S. and D. P. Burkitt (1975). Diverticular disease of the colon, a 20th Century problem. In: *Clinics of Gastroenterology*, A. N. Smith (ed.) London, Saunders, Vol. 4. p. 3.



- Patrick, G. (1967). *Nature* **216**:815. Inhibition of strontium and calcium uptake by rat duodenal slices: comparison of polyuronides and related substances.
- Pearl, I. A. (1967). *The Chemistry of Lignin*. London, Arnold.
- Ranby, B. (1969). Recent advances in the structure and morphology of cellulose. In: *Celluloses and their Applications; Advances in Chemistry Series* **95**: 245.
- Raymond, W. F. (1969). The nutritive value of forage crops. In: *Advances in Agronomy*. N. C. Brady (ed.) New York. Academic Press. Vol 21, p. 1.
- Retief, D. H. et al (1970). *Br. Dent. J.* **138**:463. Dental caries and sugar intake in South African pupils of 16 to 17 years in four ethnic groups.
- Southgate, D. A. T. and Durnin, J. V. G. A. (1970). *Br. J. Nutr.* **24**: 517. Calorie conversion factors. An experimental reassessment of the factors used in the calculation of the energy value of human diets.
- Southgate, D. A. T. (1976). In: *Fibre in Human Nutrition*. G. Spiller and R. Amen (eds.) Plenum p. 83-84.
- Thwe, M. et al (1976). Unpublished results. Quoted in: *Fibre in Human Nutrition*. G. Spiller and R. Amen (eds.) Plenum p. 200.
- van Soest, P. J. (1963). *J. Associ. Off. Agric. Chem.* **46**: 825. Use of detergents in the analysis of fibrous feeds. 1. Preparation of fibre residues of low nitrogen content.
- van Soest, P. J. (1973). *Fed Proc.* **32**: 1804. The uniformity and nutritive availability of cellulose.
- van Soest, P. J. and R. W. McQueen (1973). *Proc. Nutr. Soc.* **32**: 123. The chemistry and estimation of fibre.
- Walker, A. R. P. et al (1973). *Postgrad. Med. J.* **49**:243. Appendicitis, fibre intake and bowel behaviour in ethnic groups in South Africa.
- Wandett, S. (1969). *Nutr. Abstr. Rev.* **39**: 573. Statistical survey of the relation between sugar consumption and dental caries.
- Wattenberg, L. W. (1974). *Am. J. Dig. Dis.* **19**:947. Potential inhibitors of colon carcinogenesis.
- Wells, A. F. and B. H. Ershoff (1961). *J. Nutr.* **74**:87. Beneficial effects of pectin in prevention of hypocholesterolaemia and increase in liver cholesterol in cholesterol-fed rats.
- Wersh, S. C. and A. C. Ivy (1941). *Am. J. Dig. Dis.* **8**: 101. On the fate of ingested pectin.
- Whistler, R. L. and E. L. Richards (1970). Hemicelluloses. In: *The Carbohydrates*. W. Pigman and D. Horton (eds.) New York, Academic Press. Vol 11A p. 447.
- Williams, R. D. and Olmsted, W. H. (1936). *J. Nutr.* **11**:433. The effect of cellulose, hemicellulose and lignin on the weight of stool. A contribution to the study of laxation in man.
- Wynder, E. L. and T. Shigematsu. *Cancer* **20**:1520. Environmental factors of cancer of the colon and rectum.

# DISTRIBUTION LIST

	No. of Copies		No. of Copies
<b>DEFENCE ESTABLISHMENTS</b>		<b>DEFENCE ESTABLISHMENTS Cont'd.</b>	
Department of Defence, Russell Offices, CANBERRA, A.C.T. 2600		Headquarters Operational Command, (Staff Officer Catering), RAAF, PENRITH, N.S.W. 2750	1
(a) Secretary	1	Headquarters, Support Command (CE04/E41), RAAF, Defence Centre, 366 St. Kilda Road, MELBOURNE, Vic. 3000	
(b) Chief Defence Scientist	1	Catering Supervisor, Log Branch, HQ 1 MD, Victoria Barracks, BRISBANE, Qld. 4000	1
(c) Executive Controller, Australian Defence Scientific Service	1	Headquarters, 2 Military District, Victoria Barracks, SYDNEY, N.S.W. 2000	2
(d) Controller, Service Laboratories and Trials Division	2	Headquarters, 3 Military District, Victoria Barracks, MELBOURNE, Vic. 3000	1
(e) Superintendent, Defence Science Administration Division	1	Headquarters, 4 Military District, Kewick Barracks, ADELAIDE, S.A. 5035	1
(f) JIO (DDSTI)		51 Sup Bn, Irwin Barracks, KARRAKATTA, W.A. 6010	1
Department of Defence (Army Office), Russell Offices, CANBERRA, A.C.T. 2600		Headquarters, 6 Military District, Anglesea Barracks, HOBART, Tas. 7002	1
(a) Directorate of Army Development	1	Headquarters, 7 Military District, Larrakeyah Barracks, DARWIN, N.T. 5790	1
(b) Director of Infantry	2	Royal Military College, DUNTRON, A.C.T. 2600	
(c) Director of Catering	1	(a) The Librarian, Bridges Memorial Library	1
(d) Directorate of Army Development, C/- Scientific Adviser - Army	1	(b) The Faculty Admin Officer, Faculty of Military Studies	1
Department of Defence (Air Force Office) Russell Offices, CANBERRA, A.C.T. 2600		Commandant, Australian Staff College, FORT QUEENSCLIFFE, Vic. 3255	2
(a) Air Force Scientific Adviser	1	Commanding Officer/Chief Instructor, RAAOC Centre, Milpo, BANDIANA, Vic. 3662	2
(b) Director of Catering and Services (DCATSERV-AF)	2	Commandant, RAAF Staff College, RAAF Base, Fairbairn, CANBERRA, A.C.T. 2600	2
Department of Defence, Campbell Park Offices, CANBERRA, A.C.T. 2601		Director, Institute of Aviation Medicine, POINT COOK, RAAF, Vic. 3029	2
(a) Defence Information Services Branch	10		
(b) Defence Central Library	1		
Senior Librarian, Aeronautical Research Laboratories, P.O. Box 4331, MELBOURNE, Vic. 3001	1		
Senior Librarian, Defence Research Centre, G.P.O. Box 2151, ADELAIDE, S.A. 5001	1		
Joint Tropical Trials and Research Establishment, P.O. Box 931, INNISFAIL, Qld. 4860	1		
Directorate of Supply (Technical Services), Department of Defence (Army Office), G.P.O. Box 1932R, MELBOURNE, Vic. 3001	2		
Supply Division, HQ Logistic Command, St. James Plaza, G.P.O. Box 1932R, MELBOURNE, Vic. 3001	1		
Directorate of Army Health Services, G.P.O. Box 1932R, MELBOURNE, Vic. 3001	2		
The Director of Naval Victualling, Department of Defence (Navy Office), Victoria Barracks, St. Kilda Road, MELBOURNE, Vic. 3004	2		
Headquarters, Field Force Command, Victoria Barracks, PADDINGTON, N.S.W. 2021	3		
Headquarters, Logistic Command, St. James Plaza, G.P.O. Box 1932R, MELBOURNE, Vic. 3001	3		
Headquarters, Training Command, Victoria Barracks, PADDINGTON, N.S.W. 2021	1		
		<b>CIVILIAN ESTABLISHMENTS</b>	
		The Librarian, CSIRO, Tasmanian Regional Laboratory, Stowell Avenue, HOBART, Tas. 7000	1
		The Librarian, CSIRO Division of Food Research, P.O. Box 52, NORTH RYDE, N.S.W. 2113	1
		The Librarian, CSIRO Dairy Research Laboratory, P.O. Box 20, HIGHTT, Vic. 3190	1

## CIVILIAN ESTABLISHMENTS Cont'd.

	No. of Copies
Aust. Government Analyst, Department of Science, P.O. Box 449, WODEN, A.C.T. 2606	1
Dr. R. I. Garrod, Department of Science, 568 St. Kilda Road, MELBOURNE, Vic. 3004	1
The Central Library, Department of Health, P.O. Box 100, WODEN, A.C.T. 2606	2
Dr. R. C. Hutchinson, Derwentlaken Road, OTAGO, Tas. 7402	1
Tasmanian Collection, State Library of Tasmania, 91 Murray Street, HOBART, Tas. 7000	1
Serials Section, State Library of Tasmania, 91 Murray Street, HOBART, Tas. 7000	1
Reference Section, Hellyer Regional Library, Alexander Street, BURNIE, Tas. 7320	1
Director, Food Preservation Laboratory, Department of Primary Industries, HAMILTON, Qld. 4000	1
Head, Food School, East Sydney Technical College, Forbes Street, DARLINGHURST, N.S.W. 2010	1
Professor R. A. Edwards, Head, School of Food Technology, University of New South Wales, KENSINGTON, N.S.W. 2033	1
Preliminary Processing, National Library of Australia, CANBERRA, A.C.T. 2600	2
Serials, Biomedical Library, University of New South Wales, KENSINGTON, N.S.W. 2033	1
The Medical Library, Flinders University of South Australia, BEDFORD PARK, S.A. 5042	1

## OVERSEAS ESTABLISHMENTS

### BRITAIN

Australian Army Representative, Australia House, Strand, London, WC2 B4LA, England	1
Director of Supplies of Transport (FMV), Section 82B, Room 1235, Ministry of Defence, Empress State Building, London, SW6 1TR, England	1
Adviser in Nutrition, AMD5, (Army Dept), Ministry of Defence, Landsdowne House, Berkeley Square, London, W1, England	1
Deputy Chief Scientist (Army), SAG (A) 3b, Ministry of Defence, Main Building, Whitehall, London SW 1A 2HB, England	1
The Director, Army Personnel Research Establishment, C/- Royal Aircraft Establishment, Farnborough, Hants, England	1
The Information Officer, British Food Manufacturing Industries Research Association, Randalls Road, Leatherhead, Surrey, KT227RY, England	1

## OVERSEAS ESTABLISHMENTS Cont'd.

### BRITAIN Cont'd.

	No. of Copies
The Librarian, Food Research Institute, Colney Lane, Norwich, Norfolk NR4 7UA, England	1
Superintendent, Foods and Nutrition Division, Laboratory of the Government Chemist, Cornwall House, Stamford Street, London SE1 9NQ, England	6
The Librarian, A.R.C. Meat Research Institute, Langford, Bristol, BS18 7DY, England	1
Dr. D. J. McWeeny, Ministry of Agriculture, Fisheries and Food, Food Laboratory, Haldin House, Queen Street, Norwich, Norfolk, NR2, 4SK, England.	1
Mr. R. E. Goodman, Food Science Division, Ministry of Agriculture, Fisheries and Food, Great Westminster House, Horseferry Road, London, SW1P 2AE, England	1
The Librarian, National College of Food Technology, University of Reading, St. George Avenue, Weybridge, Surrey, England	1
The Director, Tropical Products Institute, Gray's Inn Road, London WC1X 8LU, England	1

### CANADA

National Defence Headquarters, Ottawa, Ontario, K1A 0K2	
(a) Major Jean Wallace (DCGE 3-4)	2
(b) D Food S	1
(c) DGOS Technical Library	1

### SRI LANKA

Director, Food Research and Nutrition Council, C/- Food Commissioner, Union Place, Colombo, Sri Lanka	1
--	---

### FEDERAL REPUBLIC OF GERMANY

Armed Services Food Chemist, 89 Supply Depot, RAOC (BFPO 40) Viersen, Federal Republic of Germany	1
--	---

### GHANA

Defence Adviser, Ghana High Commission, 13 Belgrave Square, London, SW1X 8PR, England	5
--	---

### INDIA

Director General, Research and Development Organisation, Ministry of Defence, New Delhi, 11	2
The Director, Defence Food Research Laboratory, Jyothi Nagar, Mysore 570 010, India	2

### MALAYSIA

Director, Defence Research Centre, Ministry of Defence, Rifle Range Road, Kuala Lumpur, Malaysia	3
---	---



**OVERSEAS ESTABLISHMENTS Cont'd.**

	No. of Copies		No. of Copies
<b>NEW ZEALAND</b>		<b>PHILIPPINES</b>	
Chief Scientist, Defence Science Organisation, DSIR, P.O. Box 8010, Wellington, New Zealand	1	Science Research Supervisor, Food Research Division, Food and Nutrition Research Institute, Manila, Philippines	1
Assistant Chief Defence Staff (Support Services) Ministry of Defence, Private Bag, Wellington, New Zealand	2		
DADST(S) Home Command, Ministry of Defence, Private Mail Bag, Wellington, New Zealand	1	<b>UNITED STATES OF AMERICA</b>	
<b>PAPUA NEW GUINEA</b>		Chief, Food and Nutrition Section, NASA — Manned Spacecraft Center, Houston, Texas	
Headquarters, Australian Defence Cooperation Group, (HQ ADCG), P.O. Box 2270, Konedobu, Port Moresby	3	Director, U.S. Army Natick Laboratories (Food Laboratory) Natick, Massachusetts, U.S.A.	2
		Director, U.S. Army Medical Research and Nutrition Laboratory, Fitzsimons General Hospital, Denver, Colorado, U.S.A. 80240	2